



**University of  
Zurich<sup>UZH</sup>**

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2014

---

## **Crohn's disease: loss of tolerance or a disorder of autophagy?**

Spalinger, Marianne R ; Rogler, Gerhard ; Scharl, Michael

**Abstract:** Crohn's disease (CD) is characterized by a breakdown of the intestinal epithelial barrier function leading to an uncontrolled immune response to bacterial antigens. Available data demonstrate that appropriate response and early host defense against invading bacteria are crucial to maintain tolerance towards commensal bacteria. When the mechanisms of early removal of invading bacteria are disturbed, a loss of tolerance and a full-blown adaptive immune reaction, which is mounted against the usually harmless commensal flora, are induced. Dysfunction of autophagy caused by genetic variations within CD susceptibility genes, such as ATG16L1 and IRGM, results in defective handling of intracellular and invading bacteria and causes prolonged survival and defective clearance of those microbes. Dysfunction of ATG16L1 and IRGM has also been shown to cause aberrant Paneth cell function and uncontrolled secretion of proinflammatory cytokines finally resulting in increased susceptibility to bacterial infection and the onset of colitis. Interestingly, autophagy can also be regulated by other CD susceptibility genes, such as NOD2 (nucleotide oligomerization domain 2) or PTPN2 (protein tyrosine phosphatase nonreceptor type 2) and the presence of the CD-associated variations within these genes results in similar effects. Taken together, more and more evidence suggests a close functional correlation between loss of tolerance and defective autophagy in CD patients. Therefore, most likely, the onset of CD is triggered by both a loss of tolerance as well as a dysfunction of autophagy, which finally results in the onset of chronic intestinal inflammation.

DOI: <https://doi.org/10.1159/000358140>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-105051>

Journal Article

Published Version

Originally published at:

Spalinger, Marianne R; Rogler, Gerhard; Scharl, Michael (2014). Crohn's disease: loss of tolerance or a disorder of autophagy? *Digestive Diseases*, 32(4):370-377.

DOI: <https://doi.org/10.1159/000358140>

# Crohn's Disease: Loss of Tolerance or a Disorder of Autophagy?

Marianne R. Spalinger<sup>a</sup> Gerhard Rogler<sup>a, b</sup> Michael Scharl<sup>a, b</sup><sup>a</sup>Division of Gastroenterology and Hepatology, University Hospital Zurich, and <sup>b</sup>Zurich Center for Integrative Human Physiology, University of Zurich, Zurich, Switzerland

## Key Words

Autophagy · Bacteria · Crohn's disease · IL-10 · Intestinal macrophages · TLR4

## Abstract

Crohn's disease (CD) is characterized by a breakdown of the intestinal epithelial barrier function leading to an uncontrolled immune response to bacterial antigens. Available data demonstrate that appropriate response and early host defense against invading bacteria are crucial to maintain tolerance towards commensal bacteria. When the mechanisms of early removal of invading bacteria are disturbed, a loss of tolerance and a full-blown adaptive immune reaction, which is mounted against the usually harmless commensal flora, are induced. Dysfunction of autophagy caused by genetic variations within CD susceptibility genes, such as ATG16L1 and IRGM, results in defective handling of intracellular and invading bacteria and causes prolonged survival and defective clearance of those microbes. Dysfunction of ATG16L1 and IRGM has also been shown to cause aberrant Paneth cell function and uncontrolled secretion of proinflammatory cytokines finally resulting in increased susceptibility to bacterial infection and the onset of colitis. Interestingly, autophagy can also be regulated by other CD susceptibility genes, such as NOD2 (nucleotide oligomerization domain 2) or

PTPN2 (protein tyrosine phosphatase nonreceptor type 2) and the presence of the CD-associated variations within these genes results in similar effects. Taken together, more and more evidence suggests a close functional correlation between loss of tolerance and defective autophagy in CD patients. Therefore, most likely, the onset of CD is triggered by both a loss of tolerance as well as a dysfunction of autophagy, which finally results in the onset of chronic intestinal inflammation.

© 2014 S. Karger AG, Basel

## Introduction

Along with ulcerative colitis (UC), Crohn's disease (CD) is one of the major forms of inflammatory bowel disease (IBD). The interplay between environmental factors, genetic predisposition, intestinal microbiota as well as the innate and adaptive immune system needs to be tightly controlled. An imbalance in these factors predisposes to the pathogenesis of IBD. Current hypotheses suggest that an epithelial barrier defect coupled with an impaired immune response of the innate as well as adaptive immune system to the commensal flora results in either excessive upregulation or impaired suppression of inflammatory responses, finally leading to chronic intes-

tinal inflammation [1]. Current studies suggested that intestinal microbial diversity is reduced and species composition is altered, especially in CD: Firmicutes are reduced while Bacteroidetes and Enterobacteriaceae are increased in the intestine of CD patients [2, 3]. Genome-wide association studies identified genetic variations in 163 genes which are associated with IBD, including 110 loci shared by both disease subtypes, 30 loci classified as specific for CD and 23 as specific for UC [4]. Of note, members of the innate immune system, such as the intracellular bacterial-sensor NOD2 (nucleotide oligomerization domain 2) as well as the autophagy genes ATG16L1 (autophagy-related gene 16-like 1) and IRGM (immunity-related GTPase family M), are exclusively associated with CD [4].

### CD – Loss of Tolerance?

Immune cells trigger an effective inflammatory response as soon as pathogen-associated molecular patterns (PAMPs) are present. In the gut, however, myriads of commensal bacteria or food antigens do normally not provoke adverse reactions. This seems to be due to active tolerance induction by intestinal epithelial cells (IECs) and specific immune cell subsets. IECs secrete mucins and antimicrobial molecules to limit the number of commensals directly penetrating the gut mucosa [5]. In CD, this barrier is impaired and increased translocation of bacteria and food antigens provokes an enhanced proinflammatory response of the innate as well as adaptive immune system, which finally results in loss of tolerance towards commensal antigens [6, 7].

#### *Intestinal Macrophages*

Myeloid intestinal immune cells, such as intestinal macrophages (iMACs) and dendritic cells (DCs) – in contrast to macrophages/DCs in other tissues, display a hypo-responsive, tolerance-inducing phenotype [8]. iMACs from CD patients, however, display a significantly higher expression of CD14, CD16, HLA-DR, CD11b, and CD11c, indicating additional macrophage populations in the inflamed mucosa, which reflects either a recruitment of new cells from the blood or a phenotypic change in resident cells [9, 10]. Under normal conditions, iMACs can eliminate intracellular bacteria without provoking an immune reaction [11]. They sample the luminal content and, by expression of specific surface markers and cytokine secretion, induce tolerogenic T cells [8]. These iMACs from normal mucosa rarely express the co-

stimulatory molecules CD80 and CD86. In CD, a macrophage population is found with high expression of costimulatory molecules, which is presumably responsible for the perpetuated immune response [12]. This special population of CD iMACs also reveals increased activation of nuclear factor (NF)- $\kappa$ B in the inflamed mucosa [13], which might, at least in part, be due to decreased levels of human glucocorticoid receptor in mucosal CD iMACs and lead to decreased protection against NF- $\kappa$ B action [14]. Further, NADPH oxidase mRNA is down-regulated or absent in macrophages from normal mucosa, which correlates with a lack of oxidative burst activity. In CD, however, macrophage oxidative burst activity is increased and NADPH oxidase mRNA is induced [15]. CD iMACs also display increased expression of TLR2 and TLR4, which likely contributes to their increased activation status in response to PAMPs. The absence of TLRs abolishes the reactivity of mucosal macrophages to bacterial wall products, while the presence of TLRs might contribute to the inflammatory process [16]. Of note, the chaperone molecule glycoprotein gp96 links the adaptive with the innate immune system. During cellular stress, peptide-loaded gp96 can be released and presented to T cells by antigen-presenting cells. gp96 is induced during differentiation of normal iMACs but is not detected in iMACs in CD mucosa. As gp96 has been described as having a role in tolerance induction, this may be relevant for loss of tolerance against luminal bacteria found in CD patients [17]. Further, a reduced expression of subunits of the ubiquitin-proteasome system in iMACs of normal mucosa supports the concept of the presence of a nonreactive, anergic macrophage phenotype in the gut under normal conditions. Reinduction in iMACs of IBD mucosa reflects activated iMACs which can present antigenic peptides and thus support inflammation [18].

### Immune Cells and Pattern Recognition Receptors

As it is the case for IECs, NOD2 malfunction in DCs and iMACs results in poor bacterial clearance [6, 19] and consequently in prolonged bacterial infection and hyperactivation of the immune system [20], finally promoting the secretion of proinflammatory cytokines and an immune-activating phenotype. NLRP3 and NLRP6 are able to form multiprotein complexes called inflammasomes, which are important for processing pro-IL-1 $\beta$  and pro-IL-18 into their mature forms and their final secretion [21]. Low-level inflammasome activation and IL-18 secretion is essential for wound healing and repair in the

intestines [21]. Physiologic NLRP3 and NLRP6 activity is involved in shaping the composition of the intestinal microbiota [22, 23]. However, uncontrolled inflammasome activity results in enhanced proinflammatory responses and proliferation of destructive immune cells [21].

Several proteins are involved in the modulation of intracellular signaling pathways downstream of pattern recognition receptors and the outcome of the resulting immune response. The CD risk gene PTPN2 (protein tyrosine phosphatase nonreceptor type 2) [24] negatively controls the activation of proinflammatory signaling cascades [25, 26]. If PTPN2 is lost or – as in some CD patients – malfunctioning, it results in hyperactivation of monocytes by bacterial products [27] and increased responses to the proinflammatory cytokines TNF and IFN- $\gamma$  [26, 28]. Finally, an inflammatory response with increased levels of IL-6, IL-12, and IFN- $\gamma$  as well as a loss of intestinal barrier function is the result, leading to the breakdown of tolerance mechanisms [26]. PTPN22 also controls bacterial-induced signaling cascades [29] and a variant within the PTPN22 gene locus protects from CD [30]. In CD, intestinal levels of PTPN22 are reduced, resulting in enhanced secretion of IL-6 but reduced IL-12 secretion [31]. This leads to a skewed response to pathogens inducing a cytokine profile favoring pathogenic Th17 cell proliferation.

At a functional level, alterations in pattern recognition receptor signaling in innate immune cells result in the aberrant induction of adaptive immune responses. During intestinal inflammation, intestinal CD14<sup>+</sup> cells secrete enhanced levels of IL-1 $\beta$ , IL-6, and IL-12 and thus promote the development of Th1 and Th17 cells [32]. Th17 cells, which defend extracellular bacteria and fungi in healthy persons [33], are increased in IBD patients and play a major role in driving pathologic inflammation and tissue destruction in the intestine [34]. In the intestinal T cell population, high proportions of so-called inducible regulatory T cells (T<sub>reg</sub>) can be found [35]. They actively downmodulate excessive activation of the adaptive immune system. In contrast to natural T<sub>reg</sub>, inducible T<sub>reg</sub>s develop in the periphery upon antigen challenge and relieve inflammatory reactions after antigen clearance following an infection [36]. In the gut, microbial-specific inducible T<sub>reg</sub>s are believed to induce tolerance towards commensal microflora [37]. In CD, however, this tolerance-inducing mechanism seems to be lost: The tight junctions between epithelial cells become leaky [7], and consequently increased numbers of commensal bacteria can penetrate the mucosa finally and thus increase NOD2 activation [6]. Further, high numbers of activated proin-

flammatory T cells can be found [38]. Additionally, IL-10 is an important regulatory cytokine able to suppress the synthesis of proinflammatory mediators, including IFN- $\gamma$  and TNF, as well as costimulatory molecules and MHC class II molecules on antigen-presenting cells [39]. T<sub>reg</sub>s and to a lower extent also Th2 cells produce IL-10 in order to downregulate proinflammatory responses and end an immune reaction restoring tolerance towards commensal bacteria [39]. The importance of this tolerance-inducing functions of IL-10 is highlighted by the fact that loss of IL-10 results in spontaneous colitis in mice [40], and loss-of-function variants in either IL-10 or IL-10 receptors result in severe, very-early-onset IBD [41].

### CD – Dysfunction of Autophagy?

Autophagy is critically involved in regulating innate immune responses by providing a defense mechanism against intracellular microbial pathogens and mediating antigen presentation via MHC class II molecules [42, 43]. Dysfunctional autophagy is associated with defective bacterial handling, prolonged intracellular survival of pathogens, and uncontrolled inflammation. Levels of autophagy proteins, such as ATG16L1 or IRGM, are significantly decreased in the intestines of CD patients [44]. To date, three autophagy genes have been confirmed as CD susceptibility genes, namely ATG16L1, IRGM, and LRRK2 (leucine-rich repeat kinase 2) [24, 45]. LRRK2-deficient mice are more susceptible to dextran sodium sulfate (DSS)-induced colitis, since LRRK2 inhibits activation of NFAT1, which promotes the secretion of proinflammatory cytokines [46]. LRRK2 expression is elevated in intestinal tissue of CD patients, and LRRK2-deficient cells are impaired in killing of intracellular bacteria [47], which suggests a possible role for LRRK2 in CD pathogenesis.

### ATG16L1, Interacting Molecules, and Paneth Cell Function

ATG16L1 represents a key molecule within the autophagy network which is responsible for subcellular localization of the autophagy machinery [48]. Presence of the SNP rs2241880 within the gene locus encoding ATG16L1 results in substitution of threonine by alanine (T300A) and has been strongly associated with an increased risk for developing CD [24]. Recent studies demonstrated that the presence of the CD-associated ATG16L1 variant results in increased numbers, pro-

longed survival, and elevated replication of intracellular bacteria [42, 43] and consequently elevated secretion of the proinflammatory cytokines TNF and IL-6, which promote inflammatory conditions in the intestine [49]. Presence of the ATG16L1 variation obviously only affects the extent of autophagy following activation, while basal levels of autophagy remain unaffected. A recent study demonstrated that CD patients featuring the ATG16L1 variation exhibit impaired antigen uptake and processing as well as a defective interaction between DCs and IECs in the intestine [50]. In a study using mice lacking the conserved coiled-coil domain of ATG16L1, increased inflammasome activation in response to PAMPs causing enhanced secretion of IL-1 $\beta$  and IL-18 was found and the mice were more susceptible to DSS-induced colitis [51]. Recent studies identified a role for microRNAs in down-regulating ATG16L1 and autophagy resulting in prolonged bacterial persistence [52, 53].

ATG16L1 seems to exert a close functional correlation with other IBD susceptibility genes, such as NOD2 or PTPN2, in regulating autophagy. In DCs, NOD2-mediated autophagy is crucial for handling of invading bacteria as well as for antigen presentation and the induction of antigen-specific CD4<sup>+</sup> T-cell responses via MHC class II molecules. DCs from CD patients carrying either the CD-associated ATG16L1 or NOD2 variations are defective in autophagosome formation, bacterial trafficking, and antigen presentation [42]. As a possible mechanism, NOD2 is crucial for the initiation of autophagy by recruiting ATG16L1 to the cell membrane at the site of bacterial entry. Cells featuring CD-associated NOD2 polymorphisms are unable to direct ATG16L1 to the plasma membrane and are deficient in handling of invading bacteria [43]. Presence of the CD-associated ATG16L1 variant resulted in increased secretion of the proinflammatory cytokines IL-1 $\beta$  and IL-6 from peripheral blood mononuclear cells of CD patients in response to stimulation with NOD2 ligands [54]. These observations strongly suggest that NOD2 as well as autophagy play a key role for the innate immune system and present the functional mechanism how dysfunction of autophagy could essentially contribute to the onset of chronic intestinal inflammation. A further close functional correlation exists between PTPN2, ATG16L1, NOD2, and autophagy in general. PTPN2 expression is, at least partially, regulated by ATG16L1 and NOD2. At a functional level, PTPN2 is involved in NOD2-mediated autophagy, since presence of the CD-associated PTPN2 variations results in impaired autophagosome formation and is associated with decreased expression of a number of autophagy genes, including IRGM and

ATG16L1 [27, 44]. Of note, a further protein tyrosine phosphatase, namely PTPN22, has also been demonstrated to regulate NOD2-mediated autophagy [29].


Besides its role in the handling of invading pathogens, ATG16L1 has been critically associated with the function of Paneth cells located within the crypts of Lieberkühn in the small intestine. Paneth cells represent a specialized epithelial cell type, which is important for the secretion of antimicrobial factors into the intestinal lumen via the production and secretion of characteristic cytoplasmic granules. In particular, hypomorphic ATG16L1 (ATG16L1<sup>HM</sup>) mice lack the antibacterial enzyme lysozyme in the mucus of ileal sections, and lysozyme was diffusely detectable in a number of Paneth cells in ATG16L1<sup>HM</sup> mice accompanied by aberrant, disorganized granules as well as decreased numbers of granules [55]. Paneth cells from ATG16L1<sup>HM</sup> mice further revealed increased expression of proinflammatory molecules, such as the adipocytokines leptin and adiponectin [55]. Of special interest, Paneth cells derived from CD patients homozygous for the ATG16L1 CD risk allele feature similar abnormalities in Paneth cell morphology and granule secretion as ATG16L1<sup>HM</sup> mice and endoplasmic reticulum stress even during quiescent CD-associated bacterial persistence in the intestine [55, 56]. Interestingly, all of the mentioned abnormalities in Paneth cell morphology and function were absent when ATG16L1<sup>HM</sup> mice were raised in an enhanced barrier facility, but could be completely introduced again following infection of the ATG16L1<sup>HM</sup> mice with the murine norovirus strain CR6 for 7 days, which indicates a critical role for a virus-plus-susceptibility gene interaction [57]. ATG16L1<sup>HM</sup> mice only displayed a severe intestinal injury response to DSS treatment when they were infected with CR6 at least 7 days before DSS administration, but not in the absence of the virus or when the virus infection occurred at the same time as the DSS treatment initiation. These observations suggest that virus-plus-susceptibility gene interactions in addition to environmental factors and commensal bacteria are critically involved in the pathogenesis of chronic intestinal inflammation.

## IRGM

Several variations within the IRGM gene locus are associated with CD, e.g. the coding region of IRGM, a 20-kb deletion polymorphism upstream of the IRGM transcriptional start site, affects several transcription factor binding sites and hereby impairs IRGM expression levels and



**Fig. 1.** CD pathogenesis: an epithelial barrier defect favors the invasion of commensal and pathogenic bacteria, food antigens, and toxins into the gut mucosa. The presence of genetic variations, such as in autophagy-related genes or pattern recognition receptors, contributes to the barrier defect and causes an impaired immune response. Hereby, alterations in tolerance mechanisms affecting the innate and adaptive immune system as well as in autophagy can be observed. Of particular interest hereby seems to be CD-specific alterations in intestinal macrophages. The aberrant immune response results in an imbalance between pro- and anti-inflammatory cytokines finally establishing the onset of chronic intestinal inflammation.



the promoter region or the 5'-untranslated region of IRGM [24, 58, 59]. Reduced expression of IRGM seems to be responsible for the decreased function of IRGM and a subsequent impairment in autophagy, which suggests that a certain level of IRGM protein is necessary for proper protein function. The necessary threshold level can obviously not be reached in the presence of the CD-associated polymorphisms and lower levels of IRGM expression have been detected in lymphocytes from CD patients [58]. Interestingly, a recent study demonstrated that the microRNA family, miR-196, which is overexpressed in the inflamed intestinal epithelium of CD patients, causes downregulation of the protective IRGM variant, but does not affect levels of the disease-associated variant. The resulting decrease in IRGM expression levels contributes to impaired autophagy and enhanced intracellular bacterial replication [60]. It has been shown that functional IRGM1 plays a key role in regulating the maturation of pathogen-containing vacuoles, as well as the adhesion and motility of activated macrophages. Consequently, IRGM1 deficiency results in increased susceptibility of mice to pathogens and finally systemic infection [61–63]. IRGM is also crucial for IFN- $\gamma$ -mediated autophagy and the elimina-

tion of intracellular *Mycobacterium tuberculosis* from human macrophages [64]. siRNA-induced knockdown of IRGM in human cells results in a defect in the autophagy machinery favoring the persistence of bacteria resulting in increased proinflammatory responses [49, 65]. Interestingly, IRGM-deficient mice featured aberrant Paneth cell morphology and localization, which is associated with impaired autophagy and increased susceptibility to DSS-induced colitis [66].

## Conclusion

Taken together, available data demonstrate that appropriate response and early host defense against invading bacteria is crucial to maintain tolerance towards commensal bacteria. If the mechanisms of early removal of invading bacteria are disturbed, this leads to a loss of tolerance and a full-blown adaptive immune reaction, which is mounted against the usually harmless commensal flora. Additionally, a number of studies clearly provide evidence for the involvement of dysfunctional autophagy in the pathogenesis of CD. Dysfunction of autophagy caused

by genetic variations within CD susceptibility genes results in defective handling of intracellular and invading bacteria as well as aberrant Paneth cell function and uncontrolled secretion of proinflammatory cytokines and thus in increased susceptibility to bacterial infection and the onset of colitis. However, more and more evidence suggests a close functional correlation between loss-of-tolerance and defective autophagy in CD patients, and all of these effects can be observed during human CD (fig. 1).

Therefore, most likely, the onset of CD is dependent on both a loss of tolerance as well as a dysfunction of autophagy, which finally results in the onset of chronic intestinal inflammation in humans.

## Disclosure Statement

The authors declare that no conflicts of interest exist.

## References

- Xavier RJ, Podolsky DK: Unravelling the pathogenesis of inflammatory bowel disease. *Nature* 2007;448:427–434.
- Walker AW, Sanderson JD, Churcher C, Parkes GC, Hudspeth BN, Rayment N, Brostoff J, Parkhill J, Dougan G, Petrovska L: High-throughput clone library analysis of the mucosa-associated microbiota reveals dysbiosis and differences between inflamed and non-inflamed regions of the intestine in inflammatory bowel disease. *BMC Microbiol* 2011;11:7.
- Sha S, Xu B, Wang X, Zhang Y, Wang H, Kong X, Zhu H, Wu K: The biodiversity and composition of the dominant fecal microbiota in patients with inflammatory bowel disease. *Diagn Microbiol Infect Dis* 2013;75:245–251.
- Jostins L, Ripke S, Weersma RK, Duerr RH, McGovern DP, Hui KY, Lee JC, Schumm LP, Sharma Y, Anderson CA, Essers J, Mitrovic M, Ning K, Cleynen I, Theatre E, Spain SL, Raychaudhuri S, Goyette P, Wei Z, Abraham C, Achkar JP, Ahmad T, Amininejad L, Ananthakrishnan AN, Andersen V, Andrews JM, Baidoo L, Balschun T, Bampton PA, Bitton A, Boucher G, Brand S, Buning C, Cohain A, Cichon S, D'Amato M, De Jong D, Devaney KL, Dubinsky M, Edwards C, Ellinghaus D, Ferguson LR, Franchimont D, Fransen K, Garray R, Georges M, Gieger C, Glas J, Haritunians T, Hart A, Hawkey C, Hedl M, Hu X, Karlsen TH, Kupcinskas L, Kugathasan S, Lattiano A, Laukens D, Lawrance IC, Lees CW, Louis E, Mahy G, Mansfield J, Morgan AR, Mowat C, Newman W, Palmieri O, Ponsioen CY, Potocnik U, Prescott NJ, Regueiro M, Rotter JI, Russell RK, Sanderson JD, Sans M, Satsangi J, Schreiber S, Simms LA, Sventoraityte J, Targan SR, Taylor KD, Tremelling M, Verspaget HW, De Vos M, Wijmenga C, Wilson DC, Winkelmann J, Xavier RJ, Zeissig S, Zhang B, Zhang CK, Zhao H, Silverberg MS, Annesse V, Hakonarson H, Brant SR, Radford-Smith G, Mathew CG, Rioux JD, Schadt EE, Daly MJ, Franke A, Parkes M, Vermeire S, Barrett JC, Cho JH: Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease. *Nature* 2012;491:119–124.
- Phillipson M, Johansson ME, Henriksnas J, Petersson J, Gendler SJ, Sandler S, Persson AE, Hansson GC, Holm L: The gastric mucus layers: constituents and regulation of accumulation. *Am J Physiol Gastrointest Liver Physiol* 2008;295:G806–G812.
- Kosovac K, Brenmoehl J, Holler E, Falk W, Schoelmerich J, Hausmann M, Rogler G: Association of the NOD2 genotype with bacterial translocation via altered cell-cell contacts in Crohn's disease patients. *Inflamm Bowel Dis* 2010;16:1311–1321.
- Zeissig S, Burgel N, Gunzel D, Richter J, Mankertz J, Wahnschaffe U, Kroesen AJ, Zeitl M, Fromm M, Schulzke JD: Changes in expression and distribution of claudin 2, 5 and 8 lead to discontinuous tight junctions and barrier dysfunction in active Crohn's disease. *Gut* 2007;56:61–72.
- Mowat AM: Anatomical basis of tolerance and immunity to intestinal antigens. *Nat Rev Immunol* 2003;3:331–341.
- Rogler G, Andus T, Aschenbrenner E, Vogl D, Falk W, Scholmerich J, Gross V: Alterations of the phenotype of colonic macrophages in inflammatory bowel disease. *Eur J Gastroenterol Hepatol* 1997;9:893–899.
- Rogler G, Hausmann M, Vogl D, Aschenbrenner E, Andus T, Falk W, Andreesen R, Scholmerich J, Gross V: Isolation and phenotypic characterization of colonic macrophages. *Clin Exp Immunol* 1998;112:205–215.
- Bilborough J, Viney JL: Gastrointestinal dendritic cells play a role in immunity, tolerance, and disease. *Gastroenterology* 2004;127:300–309.
- Rogler G, Hausmann M, Spottl T, Vogl D, Aschenbrenner E, Andus T, Falk W, Scholmerich J, Gross V: T-cell co-stimulatory molecules are upregulated on intestinal macrophages from inflammatory bowel disease mucosa. *Eur J Gastroenterol Hepatol* 1999;11:1105–1111.
- Rogler G, Brand K, Vogl D, Page S, Hofmeister R, Andus T, Knuechel R, Baeuerle PA, Scholmerich J, Gross V: Nuclear factor kappaB is activated in macrophages and epithelial cells of inflamed intestinal mucosa. *Gastroenterology* 1998;115:357–369.
- Rogler G, Meinel A, Lingauer A, Michl J, Zietz B, Gross V, Lang B, Andus T, Scholmerich J, Palitzsch KD: Glucocorticoid receptors are down-regulated in inflamed colonic mucosa but not in peripheral blood mononuclear cells from patients with inflammatory bowel disease. *Eur J Clin Invest* 1999;29:330–336.
- Hausmann M, Spottl T, Andus T, Rothe G, Falk W, Scholmerich J, Herfarth H, Rogler G: Subtractive screening reveals up-regulation of NADPH oxidase expression in Crohn's disease intestinal macrophages. *Clin Exp Immunol* 2001;125:48–55.
- Hausmann M, Kiessling S, Mestermann S, Webb G, Spottl T, Andus T, Scholmerich J, Herfarth H, Ray K, Falk W, Rogler G: Toll-like receptors 2 and 4 are up-regulated during intestinal inflammation. *Gastroenterology* 2002;122:1987–2000.
- Schreiter K, Hausmann M, Spoettl T, Strauch UG, Bataille F, Schoelmerich J, Herfarth H, Falk W, Rogler G: Glycoprotein (gp) 96 expression: induced during differentiation of intestinal macrophages but impaired in Crohn's disease. *Gut* 2005;54:935–943.
- Hetzenecker AM, Seidl MC, Kosovac K, Herfarth H, Kellermeier S, Obermeier F, Falk W, Schoelmerich J, Hausmann M, Rogler G: Downregulation of the ubiquitin-proteasome system in normal colonic macrophages and reinduction in inflammatory bowel disease. *Digestion* 2012;86:34–47.
- Biswas A, Petnicki-Ocwieja T, Kobayashi KS: NOD2: a key regulator linking microbiota to intestinal mucosal immunity. *J Mol Med (Berl)* 2012;90:15–24.
- Kim YG, Kamada N, Shaw MH, Warner N, Chen GY, Franchi L, Núñez G: The Nod2 sensor promotes intestinal pathogen eradication via the chemokine CCL2-dependent recruitment of inflammatory monocytes. *Immunity* 2011;34:769–780.
- Zaki MH, Boyd KL, Vogel P, Kastan MB, Lamkanfi M, Kanneganti TD: The NLRP3 inflammasome protects against loss of epithelial integrity and mortality during experimental colitis. *Immunity* 2010;32:379–391.

- 22 Elinav E, Strowig T, Kau AL, Henao-Mejia J, Thaiss CA, Booth CJ, Peaper DR, Bertin J, Eisenbarth SC, Gordon JI, Flavell RA: NLRP6 inflammasome regulates colonic microbial ecology and risk for colitis. *Cell* 2011;145:745–757.
- 23 Hirota SA, Ng J, Lueng A, Khajah M, Parhar K, Li Y, Lam V, Potentier MS, Ng K, Bawa M, McCafferty DM, Rioux KP, Ghosh S, Xavier RJ, Colgan SP, Tschopp J, Muruve D, MacDonald JA, Beck PL: NLRP3 inflammasome plays a key role in the regulation of intestinal homeostasis. *Inflamm Bowel Dis* 2011;17:1359–1372.
- 24 Wellcome Trust Case Control Consortium: Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 2007;447:661–678.
- 25 Scharl M, Paul G, Weber A, Jung BC, Docherty MJ, Hausmann M, Rogler G, Barrett KE, McCole DF: Protection of epithelial barrier function by the Crohn's disease associated gene protein tyrosine phosphatase N2. *Gastroenterology* 2009;137:2030–2040.e5.
- 26 Scharl M, McCole DF, Weber A, Vavricka SR, Frei P, Kellermeier S, Pesch T, Fried M, Rogler G: Protein tyrosine phosphatase N2 regulates TNF $\alpha$ -induced signalling and cytokine secretion in human intestinal epithelial cells. *Gut* 2011;60:189–197.
- 27 Scharl M, Mwinyi J, Fischbeck A, Leucht K, Eloranta JJ, Arikkat J, Pesch T, Kellermeier S, Mair A, Kullak-Ublick GA, Truninger K, Noeren F, Regula J, Gaj P, Pittet V, Mueller C, Hofmann C, Fried M, McCole DF, Rogler G: Crohn's disease-associated polymorphism within the PTPN2 gene affects muramyl-dipeptide-induced cytokine secretion and autophagy. *Inflamm Bowel Dis* 2012;18:900–912.
- 28 Scharl M, Hruz P, McCole DF: Protein tyrosine phosphatase non-receptor type 2 regulates IFN- $\gamma$ -induced cytokine signaling in THP-1 monocytes. *Inflamm Bowel Dis* 2010;16:2055–2064.
- 29 Spalinger MR, Lang S, Vavricka SR, Fried M, Rogler G, Scharl M: Protein tyrosine phosphatase non-receptor type 22 modulates NOD2-induced cytokine release and autophagy. *PLoS One* 2013;8:e72384.
- 30 Diaz-Gallo LM, Espino-Paisan L, Fransen K, Gomez-Garcia M, van Sommeren S, Cardena C, Rodrigo L, Mendoza JL, Taxonera C, Nieto A, Alcaín G, Cueto I, Lopez-Nevot MA, Bottini N, Barclay ML, Crusius JB, van Bodegraven AA, Wijmenga C, Ponsioen CY, Geary RB, Roberts RL, Weersma RK, Urcelay E, Merriman TR, Alizadeh BZ, Martin J: Differential association of two PTPN22 coding variants with Crohn's disease and ulcerative colitis. *Inflamm Bowel Dis* 2011;17:2287–2294.
- 31 Spalinger MR, Lang S, Weber A, Frei P, Fried M, Rogler G, Scharl M: Loss of protein tyrosine phosphatase nonreceptor type 22 regulates interferon- $\gamma$ -induced signaling in human monocytes. *Gastroenterology* 2013;144:978–988.e10.
- 32 Ogino T, Nishimura J, Barman S, Kayama H, Uematsu S, Okuzaki D, Osawa H, Haraguchi N, Uemura M, Hata T, Takemasa I, Mizushima T, Yamamoto H, Takeda K, Doki Y, Mori M: Increased Th17-inducing activity of CD14<sup>+</sup>CD163<sup>+</sup> low myeloid cells in intestinal lamina propria of patients with Crohn's disease. *Gastroenterology* 2013;145:1380–1391.e1.
- 33 Curtis MM, Way SS: Interleukin-17 in host defence against bacterial, mycobacterial and fungal pathogens. *Immunology* 2009;126:177–185.
- 34 Korn T, Bettelli E, Oukka M, Kuchroo VK: IL-17 and Th17 cells. *Annu Rev Immunol* 2009;27:485–517.
- 35 Hadis U, Wahl B, Schulz O, Hardtke-Wolenski M, Schippers A, Wagner N, Müller W, Sparwasser T, Förster R, Pabst O: Intestinal tolerance requires gut homing and expansion of FoxP3<sup>+</sup> regulatory T cells in the lamina propria. *Immunity* 2011;34:237–246.
- 36 Feuerer M, Hill JA, Mathis D, Benoist C: Foxp3<sup>+</sup> regulatory T cells: differentiation, specification, subphenotypes. *Nat Immunol* 2009;10:689–695.
- 37 Izcue A, Coombes JL, Powrie F: Regulatory lymphocytes and intestinal inflammation. *Annu Rev Immunol* 2009;27:313–338.
- 38 Zenewicz LA, Antov A, Flavell RA: CD4 T-cell differentiation and inflammatory bowel disease. *Trends Mol Med* 2009;15:199–207.
- 39 Moore KW, de Waal Malefyt R, Coffman RL, O'Garra A: Interleukin-10 and the interleukin-10 receptor. *Annu Rev Immunol* 2001;19:683–765.
- 40 Kuhn R, Lohler J, Rennick D, Rajewsky K, Müller W: Interleukin-10-deficient mice develop chronic enterocolitis. *Cell* 1993;75:263–274.
- 41 Kotlarz D, Beier R, Murugan D, Diestelhorst J, Jensen O, Boztug K, Pfeifer D, Kreipe H, Pfister ED, Baumann U, Puchalka J, Bohne J, Egritas O, Dalgic B, Kolho KL, Sauerbrey A, Buderus S, Güngör T, Enninger A, Koda YKL, Guariso G, Weiss B, Corbacioglu S, Socha P, Uslu N, Metin A, Wahbeh GT, Husain K, Ramadan D, Al-Herz W, Grimbacher B, Sauer M, Sykora KW, Koletzko S, Klein C: Loss of interleukin-10 signaling and infantile inflammatory bowel disease: implications for diagnosis and therapy. *Gastroenterology* 2012;143:347–355.
- 42 Cooney R, Baker J, Brain O, Danis B, Pichulik T, Allan P, Ferguson DJ, Campbell BJ, Jewell D, Simmons A: NOD2 stimulation induces autophagy in dendritic cells influencing bacterial handling and antigen presentation. *Nat Med* 2010;16:90–97.
- 43 Travassos LH, Carneiro LA, Ramjeet M, Hussey S, Kim YG, Magalhaes JG, Yuan L, Soares F, Chea E, Le Bourhis L, Boneca IG, Allaoui A, Jones NL, Nuñez G, Girardin SE, Philpott DJ: Nod1 and Nod2 direct autophagy by recruiting ATG16L1 to the plasma membrane at the site of bacterial entry. *Nat Immunol* 2010;11:55–62.
- 44 Scharl M, Wojtal KA, Becker HM, Fischbeck A, Frei P, Arikkat J, Pesch T, Kellermeier S, Boone DL, Weber A, Loessner MJ, Vavricka SR, Fried M, McCole DF, Rogler G: Protein tyrosine phosphatase nonreceptor type 2 regulates autophagosome formation in human intestinal cells. *Inflamm Bowel Dis* 2012;18:1287–1302.
- 45 Alegre-Abarrategui J, Christian H, Lufino MM, Mutihac R, Venda LL, Ansorge O, Wade-Martins R: LRRK2 regulates autophagic activity and localizes to specific membrane microdomains in a novel human genomic reporter cellular model. *Hum Mol Genet* 2009;18:4022–4034.
- 46 Liu Z, Lee J, Krummey S, Lu W, Cai H, Lenardo MJ: The kinase LRRK2 is a regulator of the transcription factor NFAT that modulates the severity of inflammatory bowel disease. *Nat Immunol* 2011;12:1063–1070.
- 47 Gardet A, Benita Y, Li C, Sands BE, Ballester I, Stevens C, Korzenik JR, Rioux JD, Daly MJ, Xavier RJ, Podolsky DK: LRRK2 is involved in the IFN- $\gamma$  response and host response to pathogens. *J Immunol* 2010;185:5577–5585.
- 48 Kuma A, Mizushima N, Ishihara N, Ohsumi Y: Formation of the approximately 350-kDa Apg12-Apg5-Apg16 multimeric complex, mediated by Apg16 oligomerization, is essential for autophagy in yeast. *J Biol Chem* 2002;277:18619–18625.
- 49 Lapaquette P, Bringer MA, Darfeuille-Michaud A: Defects in autophagy favour adherent-invasive *Escherichia coli* persistence within macrophages leading to increased pro-inflammatory response. *Cell Microbiol* 2012;14:791–807.
- 50 Strisciuglio C, Miele E, Wildenberg ME, Giugliano FP, Andreozzi M, Vitale A, Capasso F, Camarca A, Barone MV, Staiano A, Troncone R, Gianfrani C: T300A variant of autophagy ATG16L1 gene is associated with decreased antigen sampling and processing by dendritic cells in pediatric Crohn's disease. *Inflamm Bowel Dis* 2013;19:2339–2348.
- 51 Saitoh T, Fujita N, Jang MH, Uematsu S, Yang BG, Satoh T, Omori H, Noda T, Yamamoto N, Komatsu M, Tanaka K, Kawai T, Tsujimura T, Takeuchi O, Yoshimori T, Akira S: Loss of the autophagy protein Atg16L1 enhances endotoxin-induced IL-1 $\beta$  production. *Nature* 2008;456:264–268.
- 52 Lu C, Chen J, Xu HG, Zhou X, He Q, Li YL, Jiang G, Shan Y, Xue B, Zhao RX, Wang Y, Werle KD, Cui R, Liang J, Xu ZX: MIR106B and MIR93 prevent removal of bacteria from epithelial cells by disrupting ATG16L1-mediated autophagy. *Gastroenterology* 2014;146:188–199.
- 53 Zhai Z, Wu F, Chuang AY, Kwon JH: miR-106b fine tunes ATG16L1 expression and autophagic activity in intestinal epithelial HCT116 cells. *Inflamm Bowel Dis* 2013;19:2295–2301.



- 54 Plantinga TS, Crisan TO, Oosting M, van de Veerdonk FL, de Jong DJ, Philpott DJ, van der Meer JW, Girardin SE, Joosten LA, Netea MG: Crohn's disease-associated ATG16L1 polymorphism modulates pro-inflammatory cytokine responses selectively upon activation of NOD2. *Gut* 2011;60:1229–1235.
- 55 Cadwell K, Liu JY, Brown SL, Miyoshi H, Loh J, Lennerz JK, Kishi C, Kc W, Carrero JA, Hunt S, Stone CD, Brunt EM, Xavier RJ, Sleckman BP, Li E, Mizushima N, Stappenbeck TS, Virgin HW 4th: A key role for autophagy and the autophagy gene *Atg16L1* in mouse and human intestinal Paneth cells. *Nature* 2008;456:259–263.
- 56 Deuring JJ, Fuhler GM, Konstantinov SR, Peppelenbosch MP, Kuipers EJ, de Haar C, van der Woude CJ: Genomic ATG16L1 risk allele-restricted Paneth cell ER stress in quiescent Crohn's disease. *Gut* 2013, Epub ahead of print.
- 57 Cadwell K, Patel KK, Maloney NS, Liu TC, Ng AC, Storer CE, Head RD, Xavier R, Stappenbeck TS, Virgin HW: Virus-plus-susceptibility gene interaction determines Crohn's disease gene *Atg16L1* phenotypes in intestine. *Cell* 2010;141:1135–1145.
- 58 Prescott NJ, Dominy KM, Kubo M, Lewis CM, Fisher SA, Redon R, Huang N, Stranger BE, Blaszczyk K, Hudspith B, Parkes G, Hoonson N, Yamazaki K, Onnie CM, Forbes A, Dermitzakis ET, Nakamura Y, Mansfield JC, Sanderson J, Hurles ME, Roberts RG, Mathew CG: Independent and population-specific association of risk variants at the IRGM locus with Crohn's disease. *Hum Mol Genet* 2010;19:1828–1839.
- 59 McCarroll SA, Huett A, Kuballa P, Chilewski SD, Landry A, Goyette P, Zody MC, Hall JL, Brant SR, Cho JH, Duerr RH, Silverberg MS, Taylor KD, Rioux JD, Altshuler D, Daly MJ, Xavier RJ: Deletion polymorphism upstream of IRGM associated with altered IRGM expression and Crohn's disease. *Nat Genet* 2008;40:1107–1112.
- 60 Brest P, Lapaquette P, Souidi M, Lebrigand K, Cesaro A, Vouret-Craviari V, Mari B, Barbry P, Mosnier JF, Hebuterne X, Harel-Bellan A, Mograbi B, Darfeuille-Michaud A, Hofman P: A synonymous variant in IRGM alters a binding site for miR-196 and causes deregulation of IRGM-dependent xenophagy in Crohn's disease. *Nat Genet* 2011;43:242–245.
- 61 MacMicking JD, Taylor GA, McKinney JD: Immune control of tuberculosis by IFN-gamma-inducible LRG-47. *Science* 2003;302:654–659.
- 62 Henry SC, Daniell X, Indaram M, Whitesides JF, Sempowski GD, Howell D, Oliver T, Taylor GA: Impaired macrophage function underscores susceptibility to Salmonella in mice lacking *Irgm1* (LRG-47). *J Immunol* 2007;179:6963–6972.
- 63 Collazo CM, Yap GS, Sempowski GD, Lusby KC, Tessarollo L, Woude GF, Sher A, Taylor GA: Inactivation of LRG-47 and IRG-47 reveals a family of interferon gamma-inducible genes with essential, pathogen-specific roles in resistance to infection. *J Exp Med* 2001;194:181–188.
- 64 Singh SB, Davis AS, Taylor GA, Deretic V: Human IRGM induces autophagy to eliminate intracellular mycobacteria. *Science* 2006;313:1438–1441.
- 65 Lapaquette P, Glasser AL, Huett A, Xavier RJ, Darfeuille-Michaud A: Crohn's disease-associated adherent-invasive *E. coli* are selectively favoured by impaired autophagy to replicate intracellularly. *Cell Microbiol* 2010;12:99–113.
- 66 Liu B, Gulati AS, Cantillana V, Henry SC, Schmidt EA, Daniell X, Grossniklaus E, Schoenborn AA, Sartor RB, Taylor GA: *Irgm1*-deficient mice exhibit Paneth cell abnormalities and increased susceptibility to acute intestinal inflammation. *Am J Physiol Gastrointest Liver Physiol* 2013;305:G573–G584.